## AN ESTIMATE OF THE MIGRATORY TIMING OF SOCKEYE SALMON INTO UPPER COOK INLET, ALASKA, IN 1991 USING A TEST FISHERY

By

KENNETH E. TARBOX

and

BRUCE E. KING



REGIONAL INFORMATION REPORT NO. 2492-07.



Alaska Department of Fish and Game Division of Commercial Fisheries, Central Region 333 Raspberry Road Anchorage, Alaska 99581

**JUNE 1992** 

# AN ESTIMATE OF THE MIGRATORY TIMING OF SOCKEYE SALMON INTO UPPER COOK INLET, ALASKA, IN 1991 USING A TEST FISHERY

By: Kenneth E. Tarbox and Bruce E. King

Regional Information Report' No. 2A92-07

Alaska Department Of Fish And Game Division Of Commercial Fisheries 333 Raspberry Road Anchorage, Alaska 99518

June 1992

The Regional Information Report Series was established in 1987 to provide an information access system for all unpublished divisional reports. These reports frequently serve diverse ad hoc informational purposes or archive basic uninterpreted data. To accommodate timely reporting of recently collected information, reports in this series undergo only limited internal review and may contain preliminary data; this information may be subsequently finalized and published in the formal literature. Consequently, these reports should not be cited without prior approval of the author or the Division of Commercial Fisheries.

## TABLE OF CONTENTS

	<u>Page</u>
LIST OF TABLES	ii
LIST OF FIGURES	iii
LIST OF APPENDICES	iv
ABSTRACT	1
INTRODUCTION	2
METHODS	2
Test Fishing	2
Describing the Salmon Migration	3
RESULTS	4
DISCUSSION	6
LITERATURE CITED	7
APPENDICES	22

## LIST OF TABLES

Table		Page
1.	Summary of fishing time, daily and cumulative sockeye salmon catch CPUE, Upper Cook Inlet offshore test fishing project, 1991	8
2.	Estimated sockeye salmon catch by date and station, Upper Cook Inlet offshore test fishing project, 1991	9
3.	Estimated sockeye salmon CPUE by date and station, Upper Cook Inlet offshore test fishing project, 1991	10
4.	Mean date of the sockeye salmon run across Anchor Point transect by year, Upper Cook Inlet offshore test fishing project	11
5.	The 1991 Upper Cook Inlet commercial salmon harvest	12
6.	Total offshore test fishing CPUE and total sockeye salmon run estimates, Upper Cook Inlet, 1991	14
7.	Performance of 1983 run timing model as a predictor of total sockeye salmon run size in 1991, Upper Cook Inlet, offshore test fishing project	15
8.	Estimates of sockeye salmon catchability, passage rate, and relative error of estimates, Upper Cook Inlet, Alaska offshore test fishing project, 1991	16
9.	Summary of sockeye salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE at Station 6a, Upper Cook Inlet offshore test fishing project, 1991	17

## LIST OF FIGURES

<u>Figur</u>	<u>e</u>	<u>Page</u>
1.	Map of Upper Cook Inlet, Alaska, showing location of Offshore Test Fish transect	18
2.	Daily (upper) and cumulative (lower) CPUE for sockeye salmon recorded during the 1991 offshore test fish project, Upper Cook Inlet, Alaska	19
3.	Cumulative proportions estimated for the sockeye salmon return to Upper Cook Inlet, Alaska, in 1991	20
4.	Cumulative proportions for all stations combined, station 6a only, and all stations except 6a for the sockeye salmon return to Upper Cook Inlet, Alaska, in 1991	21

## LIST OF APPENDICES

<u>Table</u>		Page
A.1 -	Summary of pink salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fishing project, 1991	22
A.2 -	Estimated pink salmon catch by date and station, Upper Cook Inlet offshore test fishing project, 1991	23
A.3 -	Estimated pink salmon CPUE by date and station, Upper Cook Inlet offshore test fishing project, 1991	24
B.1 -	Summary of chum salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fishing project, 1991	25
B.2 -	Estimated chum salmon catch by date and station, Upper Cook Inlet offshore test fishing project, 1991	26
B.3 -	Estimated chum salmon CPUE by date and station, Upper Cook Inlet offshore test fishing project, 1991	27
C.1 -	Summary of coho salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fishing project, 1991	28
C.2 -	Estimated coho salmon catch by date and station, Upper Cook Inlet offshore test fishing project, 1991	29
C.3 -	Estimated coho salmon CPUE by date and station, Upper Cook Inlet offshore test fishing project, 1991	30
D.1 -	Summary of chinook salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fishing project, 1991	31
D.2 -	Estimated chinook salmon catch by date and station, Upper Cook Inlet offshore test fishing project, 1991	32

## LIST OF APPENDICES (continued)

<u>Table</u>		Page
D.3 -	Estimated chinook salmon CPUE by date and station, Upper Cook Inlet offshore test fishing project, 1991	33
E.1 -	Entry pattern of sockeye salmon into Upper Cook Inlet, Alaska, 1991, estimated from daily CPUE measured at the latitude of Anchor Point	34
F.1 -	Environmental measurements and observations made in Upper Cook Inlet, 1991, Alaska during the offshore test fishing project	35

#### **ABSTRACT**

During the 1991 Upper Cook Inlet, Alaska commercial salmon fishing season a test fishery was used to estimate the timing of the sockeye salmon, *Oncorhynchus nerka*, run as it passed a transect along the southern boundary of the management area. The test fishery operated from 1 July to 30 July and captured 2,401 sockeye salmon representing 1,574.1 CPUE points. Mean date of the run was 17 July, and the test fishery encompassed approximately 89.5% of the total run.

KEY WORDS: Salmon, Oncorhynchus, Upper Cook Inlet, Alaska, test fishery,

migratory behavior

#### INTRODUCTION

In 1979, the Alaska Department of Fish and Game (ADF&G) began a test fishing project near the southern boundary of the Upper Cook Inlet (UCI) salmon management area (Figure 1). The objective of this project was to estimate the total run of sockeye salmon, *Oncorhynchus nerka*, to UCI before salmon reached commercial harvest areas. Such information has helped ADF&G management biologists set commercial fishing times and areas to harvest sockeye salmon surplus to spawning needs. Test fishing results have been reported annually since 1979 (Waltemyer 1983a, 1983b, 1986a, 1986b, Hilsinger and Waltemyer 1987, Hilsinger 1988, Tarbox and Waltemyer 1989, Tarbox, 1990). This report presents the results of the 1991 test fishing project.

#### METHODS

#### Test Fishing

Sockeye salmon returning to Upper Cook Inlet were sampled by fishing geographically fixed stations between Anchor Point and Red River Delta (Figure 1). Stations were numbered consecutively from east to west. Station locations were determined from LORAN C coordinates. A chartered test fishing vessel sampled stations 4 - 8 daily. To determine whether an increase in sampling power would improve estimates, an additional station (6a) was sampled every other day. Data for this station were not incorporated into actual estimates but were evaluated independently for this report.

Sampling started on 1 July 91 and continued through 30 July. The chartered vessel, F/V Corrina Kay, fished 200 fm (366 m) of 2.1 cm (5 1/8 in) multifilament gill net during test fishing. Drift gill net web had a filament size number of 53/S6F, was 45 meshes deep, and was constructed of double knot Super Crystal shade number 1.

All salmon captured were identified to species. All sockeye salmon were measured for length (mid-eye to fork of tail in mm). The number of each species caught at each station was expressed as a catch per unit of effort (CPUE) statistic:

where: CPUE = CPUE for station s, and MFT = mean fishing time.

Mean fishing time was calculated as:

$$MFT = (C - B) + \frac{[B - A] + [D - C]}{2}$$

where: A = time net deployment started,

B = time net fully deployed,

C = time net retrieval started, and

D = time net fully retrieved.

Once deployed at a station, gill nets were fished 30 min before retrieval started.

Daily CPUE (CPUE<sub>a</sub>) was calculated as:

$$CPUE_s = \sum_{s=1}^{n} CPUE_s$$

The following physical and chemical measurements were taken at the start of each set: air temperature, water temperature (at 1 m below the surface), wind velocity and direction, tide stage, water depth, and water clarity. Air and water temperatures were measured using a YSI salinity/temperature meter. Unfortunately, the salinity meter malfunctioned during the project. Wind speed was measured in knots and direction was recorded as 0 (no wind), 1 (north), 2 (northeast), 3 (east), 4 (southeast), 5 (south), 6 (southwest), 7 (west), or 8 (northwest). Tide stage was classed as flood, ebb or slack by observing the movement of the vessel while drifting with the gill net. Water depth was measured in fm using a Simrad echo sounder, and water clarity was measured in m using a 17.5 cm secchi disk.

## Describing the Salmon Migration

Catchability, the fraction of the available population taken by a defined unit of fishing effort, was estimated as:

$$q_a = c_a/r_a$$
 ,

where:  $q_a$  = estimated catchability on day d,

 $r_a$  = adjusted cumulative total return on day d, and

 $c_d$  = cumulative CPUE on day d.

Passage rate, the expansion factor used to convert CPUE into estimated numbers of salmon passing the test fishing transect, was calculated as:

$$PR = 1/q_a = passage rate$$

Since the test fishery did not encompass the entire sockeye salmon run, the total CPUE for the test fishery was estimated after the season using the following relationship:

$$CPUE_{t} = CPUE_{t} \times \underbrace{H_{t}}_{H_{(t+2)}},$$

where: CPUE, = total estimated CPUE for the season,

CPUE, = cumulative CPUE through final day, f, of test fishing,

H. = total commercial harvest for the season

 $H_{(r-2)}$  = total commercial catch through final day of test fishery

(f+2), and

2 = number of days it took salmon to travel from test fishery to commercial harvest areas.

Estimates of CPUE, and CPUE, values were used to estimate daily and cumulative proportions of CPUE, based on a non-linear model:

$$y_a = 1/(1 + e^{-(a+bd)})$$

where:  $y_a = \text{cumulative proportion of CPUE or return on day d},$  a and b = coefficients of model, d = day of observation.

To calculate mean date of return, the following formula was used:

$$M = a/b$$

where: M = mean date of return,

#### **RESULTS**

A total of 2,401 sockeye salmon, 35 pink salmon, 0. gorbuscha, 671 chum salmon, 0. keta, 690 coho salmon, 0. kisutch, and 2 chinook salmon, 0. tshawytscha, were

captured during the 1991 test fishery (Table 1, Appendices A-D). Daily sockeye salmon catches ranged from 6 to 455 fish (Table 1).

Sockeye salmon daily CPUE values ranged from 5.0 on 24 July to 232.4 on 15 July. Cumulative total CPUE for the duration of the project was 1,574.1 (Table 1, Figure 2 and 3). Using post season commercial harvest figures, test fishing spanned approximately 89.5% of the total run. Therefore, total CPUE for the test fishery would have been 1,759, if test fishing had continued throughout the duration of the run.

Sockeye salmon catches along the transect were similar to the distribution of CPUE values. Approximately 86% of the total sockeye salmon catch and 84.1% of the total CPUE occurred at stations 5, 6, and 7 (Tables 2 and 3).

Examination of daily and cumulative proportions (estimated post season) of the sockeye salmon run to UCI suggested that only 1.4% of the run had passed the transect prior to the start of test fishing on 1 July and that the run was 91.6% completed at project termination (Appendix E; Figure 4). The mean date of the run was 17 July 1991 which was 3 d late relative to the historic average (Table 4).

The total sockeye salmon run to UCI in 1991 was estimated to be 3,600,000 fish of which 2,177,576 were harvested in the commercial fishery (Table 5). Estimated passage rate for the season was 2,047 sockeye salmon per CPUE index point.

Water temperatures measured along the transect generally increased during the season from a low of  $8.0^{\circ}\text{C}$  early in July to a high of  $11.5^{\circ}\text{C}$  in mid-July (Appendix F). Air temperatures fluctuated between  $8^{\circ}\text{C}$  and  $15^{\circ}\text{C}$  during the project (Appendix F). Wind velocities were generally moderate. However, winds of 20 knots or greater were recorded on 2 days (Appendix F). Wind direction was typically from the southeast or northeast.

During the commercial salmon fishing season five estimates of the sockeye salmon total run were made (Table 6). Past studies suggested that the initial best fit estimate was not accurate in predicting total run during the season and that the second or third best fit estimate was usually more useful. This pattern was again evident in 1991 as the initial best fit estimate (1990 run timing) on 11 July was 9,700,000 fish, while the third best fit (1983 run timing) was 4,200,000 fish. Although both estimators produced smaller run size predictions as the season progressed, 1990 run timing suggested the run was well above its actual size (3,600,000) until 30 July. Estimates based on 1983 run timing ranged from 3,400,000 to 4,200,000 fish. In contrast, estimates based on 1990 run timing ranged from 9,700,000 to 3,200,000 fish. During a critical management time period (16 - 18 July) the 1990 timing data suggested that the 1991 run could be 5,300,000 fish. The 1983 comparison bracketed the actual total return with a range of 3.4 to 4.1 million fish (Table 6 and 7). Consistent with previous investigations initial estimates of total CPUE were low (Table 7) and passage rate was estimated high (Table 8).

Addition of sampling effort at station 6a (Table 9) did not alter estimated mean timing of the sockeye salmon run. Visual comparison of run timing curves with and without station 6a indicated no major differences (Figure 4). Using only station 6a data to define run timing indicated that the run was 1 day earlier than was shown by combining data from all stations.

#### DISCUSSION

The 1991 season again demonstrated the continuing problem of the Upper Cook Inlet offshore test fish program. Initial estimates show a bias towards estimating timing to be later than the actual run. Therefore, estimates are generally greater than the actual return early in the season. In 1991, fishery managers in Upper Cook Inlet recognized this problem and chose to use the third best fit year as a more reasonable approximation of the run. While this proved successful for the 1991 season it was a subjective decision which may not be applicable all years. To improve the reliability of the offshore test fishing program, more effort and monies must be allocated to determine the cause of these problems. Increased sampling power using hydroacoustic techniques or more fishing vessels may be one solution.

#### LITERATURE CITED

- Hilsinger, J.R. 1988. Run strength analysis of the 1987 sockeye salmon return to Upper Cook Inlet, Alaska. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A88-19, Anchorage.
- Hilsinger, J.R. and D. Waltemyer. 1987. Run strength analysis of the 1986 sockeye salmon return to Upper Cook Inlet, Alaska. Alaska Department of Fish and Game, Division of Commercial Fisheries, Upper Cook Inlet Area Data Report 87-6, Soldotna.
- Tarbox, K.E. 1990. An estimate of the migratory timing of sockeye salmon into Upper Cook, Alaska, in 1989 using a test fishery. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2S90-4, Anchorage.
- Tarbox, K.E. and D. Waltemyer. 1989. An estimate of the 1988 total sockeye salmon return to Upper Cook Inlet, Alaska using a test fishery. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2S89-4, Anchorage.
- Waltemyer, D.L. 1983a. Migratory timing and abundance estimation of the 1982 sockeye salmon return to Upper Cook Inlet based on a test fishing program. Alaska Department of Fish and Game, Division of Commercial Fisheries, Upper Cook Inlet Data Report 83-1, Soldotna.
- 1983b. Describing the migrations of salmon and estimating abundance of sockeye salmon returning in 1983 to Upper Cook Inlet based on a test fishery. Alaska Department of Fish and Game, Division of Commercial Fisheries, Upper Cook Inlet Data Report 84-1, Soldotna.
- \_\_\_\_\_\_1986a. Use of a test fishery to describe and estimate the sockeye salmon total return to Upper Cook Inlet in 1984. Alaska Department of Fish and Game, Division of Commercial Fisheries, Upper Cook Inlet Data Report 86-1, Soldotna.
- \_\_\_\_\_\_1986b. Run strength analysis of the 1985 sockeye salmon return to Upper Cook Inlet, Alaska based on a test fishery. Alaska Department of Fish and Game, Division of Commercial Fisheries, Upper Cook Inlet Data Report 86-5, Soldotna.

Table 1. Summary of fishing time, daily and cumulative sockeye salmon catch and CPUE, Upper Cook Inlet offshore test fishing project, 1991.

	Number	Mean Fishing	Cat	ch	CF	PUE	Mean
Date	of Stations	Time (min)	Daily	Cum	Daily	Cum	Length (mm)
01 Jul 02 Jul	y 4	195.5 131.5	50 16	50 66	35.2 12.0	35.2 47.2	539 510
03 Juli 04 Juli	_	186.0 188.5	8 26	74 100	5.6 20.3	52.8 73.1	525 515
05 Jul	y 5	186.0	33	133	26.2	99.3	524
06 Juli 07 Juli	y 5	190.5 188.5	46 43	179 222	35.7 34.2	135.0 169.2	522 524
08 Jul	y 5	186.5 189.5	34	256	26.9	196.1	529 547
09 Juli 10 Juli	y 5 y 5	189.5	62 35	318 353	48.2 26.7	244.2 270.9	547 522
11 Jul 12 Jul	y 5	204.5 188.5	163 76	516 592	108.1 55.7	379.0 434.7	545 546
13 Jul	y 5 y 5	217.5	234	826	146.4	581.2	558
14 Juli 15 Juli	y 5 v 5	180.0 230.0	34 455	860 1315	25.1 232.4	606.3 838.7	556 564
16 Jul	y 5	195.0	97	1412	71.6	910.3	556
17 Juli 18 Juli	y 5 v 5	187.5 196.0	20 136	1432 1568	15.7 96.8	926.0 1022.8	557 562
19 Jul	ý 5	192.5	50	1618	37.9	1060.8	551
20 Juli 21 Juli	y 5 y 5	180.0 200.0	117 50	1735 1785	75.9 35.8	1136.7 1172.5	550 564
22 Jul		200.0 194.5	125	1910	80.8	1253.3	573 540
23 Juli 24 Juli	y 5 y 5	178.0	61 6	1971 1977	41.7 5.0	1295.0 1300.1	548 555
25 Jul 26 Jul	y 5	191.5 192.5	52 37	2029 2066	37.3 25.9	1337.4 1363.3	586
27 Jul	y 5	191.5	141	2207	84.4	1447.7	565
28 Jul 29 Jul		110.5 202.6	28 152	2235 2387	21.1 93.5	1468.8 1562.3	557 565
30 Jul		157.5	14	2401	11.8	1574.1	572

Table 2. Estimated sockeye salmon catch by date and station, Upper Cook Inlet offshore test fishing project, 1991.

			Station N	lumber		
Date	4	5	6	7	8	Total
01 July 02 July	1 0	41	6 0	0 15	2	50 16
03 July	1	3	1	2	· 1	8
04 July	13	10	0	2	1	26
05 July	20	12	0	1	0	33
06 July 07 July	2 13	7 18	14 1	7 11	16 0	46 43
07 July 08 July	4	4	19	7	0	34
09 July	2	27	12	21	Ŏ	62
10 July	0	33	0 /	1	1	35
11 July	3	90	58	3	9	163
12 July 13 July	39	36	1 48	0 87	0 2	76 234
13 July 14 July	2 3	95 0	25	5	1	34
15 July	Õ	13	241	201	Ō	455
16 July	3	20	4	45	25	97
17 July	6	6	6	0	2	20
18 July 19 July	7 16	71 29	38 1	11 2	9 2	136 50
19 July 20 July	13	100	2	1	1	117
21 July	3	12	20	15	Ō	50
22 July	0	15	17	90	3	125
23 July	1	0	9	2	49	61
24 July 25 July	0 1	0 0	3 3	3 9	0 39	6 <b>5</b> 2
25 July 26 July	0	2	24	10	1	37
27 July	ŏ	5	100	34	2	141
28 July			20	6	2	28
29 July	1	90	13	45	3	152
30 July	0	0	11	0	3	14
Total	154	740	697	636	174	2401
%	6.4	30.8	29.0	26.5	7.2	100.0

Table 3. Estimated sockeye salmon CPUE by date and station, Upper Cook Inlet offshore test fishing project, 1991.

			Station N	umber		
Date	4	5	6	7	8	Total
01 July 02 July 03 July 04 July 05 July 06 July 07 July 08 July 10 July 11 July 12 July 13 July 14 July 15 July 16 July 17 July 18 July 19 July 20 July 21 July 22 July 23 July 24 July 25 July 26 July 27 July 28 July 29 July 30 July	0.7 0.0 0.8 10.1 15.6 1.6 10.3 3.2 1.6 0.0 5.0 28.2 1.6 2.5 0.0 2.5 4.7 5.7 12.5 10.3 2.5 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0	28.0 0.8 1.6 7.7 9.7 5.9 14.6 3.2 20.5 25.0 55.7 26.7 60.0 10.8 15.4 4.8 47.4 21.5 58.3 9.6 12.2 0.0 0.0 1.7 3.3 54.5 0.0	4.6 0.0 0.8 0.0 0.0 10.8 0.8 14.8 9.5 0.0 37.4 0.8 32.7 18.1 117.6 3.2 4.5 27.1 0.8 5.7 14.0 12.7 7.0 2.5 2.5 15.5 56.3 12.0 9.3	0.0 11.2 1.6 0.8 5.6 5.6 16.8 2.5 0.4 31.8 0.0 9.6 9.8 53.5 6.8 7.2 23.2 7.5 26.5 0.0	1.8 0.8 0.0 11.9 0.0 0.0 0.0 0.8 7.5 0.0 1.6 0.8 0.0 18.7 7.6 1.6 0.8 0.0 2.4 32.3 0.0 27.2 0.8 1.6 2.4 2.5	35.2 12.0 5.6 20.3 26.2 35.7 34.2 26.9 48.2 26.7 108.1 55.7 146.4 25.1 232.4 71.6 15.7 96.8 37.9 75.9 35.8 80.8 41.7 5.0 37.3 25.9 84.4 21.1 93.5 11.8
Total	122.1	499.0	430.0	395.51	27.5	1574.1
, %	7.8	31.7	27.3	25.1	8.1	100.0

Table 4. Mean date of the sockeye salmon run across Anchor Point transect, Upper Cook Inlet offshore test fishing project, 1979-1991.

Year	Mea	n Date
	Coded	Calendar
1979	18.4	July 11
1980	22.7	July 15
1981	13.2	July 06
1982	24.2	July 17
1983	22.6	July 15
1984	18.4	July 11
1985	22.7	July 15
1986	23.0	July 16
1987	25.7	July 18
1988	20.6	July 13
1989	21.6	July 14
1990	25.6	July 18
1991	24.3	July 17
1979-1990	21.5	July 14

 $<sup>^{\</sup>circ}$  Day (1) = June 24.

Table 5. The 1991 Upper Cook Inlet commercial salmon harvest.

		Ch	inook	9	ockeye	(	Coho	Р	ink	(	Chum		Total
Date	Number of Deliveries	Daily	Cum	Daily	Cum	Daily	Cum	Daily	Cum	Daily	Cum	Daily	Cum
5-27 Mon	. 8	66	66	91	91							157	157
5-31 Fri	14	71	137	551	642							622	779
6-03 Mon		3,047	3,184	1,854	2,496							4,901	5,680
6-07 Fri		239	3,423	3,243	5,739							3,482	9,162
6-10 Mon	137	1,929	5,352	4,224	9,963							6,153	15,315
6-14 Fri	14	26	5,378	2,124	12,087					`		2,150	17,465
6-17 Mon		1,610	6,988	4,303	16,390							5,913	23,378
6-21 Fri		57	7,045	959	17,349							1,016	24,394
6~24 Mon		386	7,431	1,477	18,826	3	3	1	1	1	1	1,868	26,262
6-28 Fri	110	343	7,774	6,476	25,302	4	7	5	6	40	41	6,868	33,130
7-01 Mon	283	455	8,229	37,360	62,662	69	76	43	49	284	325	38,211	71,341
7-05 Fri	308	500	8,729	41,257	103,919	404	480	148	197	312	637	42,621	113,962
7-06 Sat		240	8,969	15,178	119,097	21	501	110	307	10	647	15,559	129,521
7-08 Mon	912	649	9,618	198,501	317,598	5,196	5,697	400	707	6,838	7,485	211,584	341,105
7-12 Fri	588	661	10,279	40,701	358,299	1,113	6,810	626	1,333	141	7,626	43,242	384,347
7-15 Mon	1107	840	11,119	556,641	914,940	47,510	54,320	3,172	4,505	26,925	34,551	635,088	1,019,435
7-18 Thu		139	11,258	118,691	1,033,631	118	54,438	65	4,570	6	34,557	119,019	1,138,454
7-19 Fri	1329	632	11,890	590,619	1,624,250	52,810	107,248	3,975	8,545	39,153	73,710	687,189	1,825,643
7-20 Sat	204	263	12,153	50,383	1,674,633	184	107,432	352	8,897	9	73,719	51,191	1,876,834
7-21 Sun	31		12,153	5,319	1,679,952	943	108,375	11	8,908	427	74,146	6,700	1,883,534
7-22 Mon	196	21	12,174	27,529	1,707,481	18,002	126,377	2,000	10,908	2,782	76,928	50,334	1,933,868
7-23 Tue	22		12,174	5,140	1,712,621	687	127,064	10	10,918	534	77,462	6,371	1,940,239
7-26 Fri			12,174		1,712,699	64	127,128	22	10,940	373	77,835	537	1,940,776
7-29 Mon	954	347	12,521	142,753	1,855,452	22,998	150,126	981	11,921	31,769	109.604	198,848	
7-31 Wed	366	129	12,650	27,994	1,883,446	5,712	155,838	119	12,040	1,647	111,251		2,175,225
8-01 Thu	569	163	12,813	65,478	1,948,924	9,198	165,036	263	12,303	5,397	116,648	80,499	2,255,724
8-02 Fri	954	194	13,007	95,132	2,044,056	58,335	223,371	1,043	13,346	43,230	159,878	197,934	2,453,658
8-03 Sat	452	133	13,140	20,405	2,064,461	4,669	228,040	132	13,478	392	160,270		2,479,389
8-04 Sun		120	13,260	17,310	2,081,771	6,382	234,422	61	13,539	398	160,668		2,503,660
8-05 Mon	862	149	13,409		2,129,685	46,113	280,535	576	14,115	52,358	213,026		2,650,770

<sup>-</sup> Continued -

Table 5. (p. 2 of 2)

		Ch	inook	S	Sockeye	(	Coho	Р	ink	(	Chum		Total
Date	Number of Deliveries	Daily	Cum	Daily	Cum	Daily	Cum	Daily	Cum	Daily	Cum	Daily	Cum
8-09 Fri	571	61	13.470	19,511	2.149.196	35,036	315,571	178	14.293	27,100	240.126	81.886	2.732.656
8-12 Mon		49	13,519	15,399	2.164.595	30,758	346,329	205	14,498	19.428	259.554	65,839	2,798,495
8-16 Fri	191	3	13,522	3,195		16,430	362,759	59	14,557	8,772	268,326	28,459	2.826.954
8-19 Mon	134	4	13,526	2,433	2,170,223	12,511	375,270	48	14,605	4,695	273,021	19,691	2.846.645
8-21 Wed	103	2	13,528	2,218	2,172,441	12,567	387,837	32	14,637	2,728	275,749	17,547	
8-23 Fri	130	2	13,530	2,184	2,174,625	11,276	399,113	19	14,656	2,402	278,151	15,883	2,880,075
8-26 Mon	74		13,530	1,131	2,175,756	7,233	406,346	2	14,658	1,473	279,624	9.839	2,889,914
8-28 Wed	47	3	13,533	1,077	2,176,833	4,333	410,679	5	14,663	188	279,812	5,606	2,895,520
8-30 Fri	40	1	13,534	264	2,177,097	3,971	414,650		14,663	294	280,106	4,530	2,900,050
9-02 Mon	44		13,534	258	2,177,355	3,356	418,006		14,663	21	280,127	3,635	2,903,685
9-04 Wed	29		13,534	57	2,177,412	2,772	420,778		14,663	22	280,149	2,851	2,906,536
9~06 Fri			13,534	124	2,177,536	2,622	423,400		14,663	17	280,166	2,763	2,909,299
9-09 Mon			13,534	5	2,177,541	690	424,090		14,663	8	280,174	703	2,910,002
9-11 Wed		1	13,535		2,177,554	740	424,830		14,663	38	280,212	792	2,910,794
9-13 Fri	12		13,535	17	2,177,571	526	425,356		14,663	9	280,221	552	2,911,346
9-16 Mon	4		13,535		2,177,571	224	425,580		14,663	2	280,223	226	2,911,572
9-18 <b>W</b> ed	4		13,535	4	2,177,575	119	425,699		14,663		280,223	123	2,911,695
9-23 Mon	1		13,535	1	2,177,576	25	425,724		14,663		280,223	26	

Table 6. Total offshore test fishing CPUE and total sockeye salmon run estimates, Upper Cook Inlet, 1991.

Date	Cum CPUE³	Best Fit Year	MSSDEV°	Passage Rate	Total CPUE	Total Run Estimate (X 10°)	Percent Error°
ll July	379.0	1990 1983	.000018	4,212 4,212	2.311	9.7 4.2	169.4 16.7
16 July	910.3	1990 1983	.000122 .002632	2,834 2,834	2,536 1,459	7.1 4.1	97.2 16.7
18 July	1022.8	1990 1983	.000198 .002889	2,160 2,160	2,473 1,563	5.3 3.4	47.2 -5.6
20 July	1136.7	1990 1983	.000553 .002688	2,454 2,454	2,326 1,596	5.7 3.9	58.3 8.3
30 July	1574.1	1990 1983	.000769	1,997 1,997	1,600 1,856	3.2 3.7	-11.1

 $<sup>\</sup>mbox{^{\circ}}$  Cumulative CPUE from start of test fishing to estimated end of total run.  $\mbox{^{\circ}}$  Mean sum of squared deviation.

Percent error = predicted-actual actual

Table 7. Performance of 1983 run timing model as a predictor of total sockeye salmon run size in 1991, Upper Cook Inlet, offshore test fishing project.

		CPUE <sup>a</sup>	Total Return°		
Date	Percent of Total Run	Estimate of Total	Percent Error <sup>c</sup> (%)	Estimate (millions)	Percent Error <sup>e</sup> (%)
11 July	23.3	1,017	-42.2	4.2	16.7
16 July	43.9	1,459	-17.1	4.1	13.9
18 July	53.3	1,563	-11.1	3.4	-5.6
20 July	62.4	1,596	- 9.3	3.9	8.3
30 July	91.6	1,856	5.5	3.7	2.8

<sup>\*</sup> Total CPUE for season was 1,759. Percent migration based on curve fit of daily index points.

<sup>\*</sup> Total sockeye salmon return was 3.6 million fish.

\* Mean percent error (MPE) = -14.8 Absolute MPE (AMPE)

<sup>&</sup>quot; MPE = 7.2 AMPE = 9.5

Table 8. Estimates of sockeye salmon catchability, passage rate, and relative error of estimates, Upper Cook Inlet, Alaska offshore test fishing project, 1991.

	Catchal	oility*	Passage Rate <sup>™</sup>		
Date	q	Percent Error (%)	q	Percent Error (%)	
ll July	0.000237	-51.5	4,212	105.8	
16 July	0.000353	-27.8	2,834	38.4	
18 July	0.000463	- 5.3	2,160	5.5	
20 July	0.000407	-16.8	2,454	19.9	
30 July	0.000500	2.2	1,997	-2.4	

<sup>&#</sup>x27;Catchabilty (q) for 1991 was 0.000489.
'Passage rate for 1991 was 2,047 fish per unit of effort?

 $<sup>^{\</sup>circ}$  MPE = 19.8

AMPE = 20.7

<sup>&</sup>quot; MPE = 33.4 AMPE = 34.4

Table 9. Summary of sockeye salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE at Station 6a, Upper Cook Inlet offshore test fishing project, 1991.

	Number	Mean Fishing	Cat	ch	СР	UE
Date	of Stations	Time (min)	Daily	Cum	Daily	Cum
02 July 03 July 04 July 05 July 06 July 07 July 08 July 10 July 11 July 12 July 13 July 14 July 15 July 16 July 17 July 18 July	1 1 0 1 0 1 0 1 0 1 0 1 0 1	40.5 36.5 .0 37.5 .0 38.5 .0 32.5 .0 52.0 .0 41.5 .0 12.0 .0 37.5	20 0 0 3 0 5 0 31 0 131 0 36 0 17 0 2 0	20 20 20 23 23 28 28 59 59 190 190 226 243 243 245 245 246	14.8 0.0 0.0 2.4 0.0 3.9 0.0 28.6 0.0 75.6 0.0 26.0 0.0 87.0 0.0	14.8 14.8 14.8 17.2 17.2 21.1 21.1 49.7 49.7 125.3 125.3 151.4 238.4 238.4 240.0 240.0 240.0
20 July 21 July 22 July 23 July 24 July 25 July 26 July 27 July 28 July 29 July	0 1 0 1 0 1 0 1	.0 48.0 .0 37.0 .0 45.5 .0 41.5 .0	0 32 0 2 0 69 0 14 0 54	246 278 278 280 280 349 349 363 363 417	0.0 20.0 0.0 1.6 0.0 45.5 0.0 13.3 0.0 34.8	240.7 260.7 260.7 262.4 262.4 307.8 307.8 321.2 356.0




Figure 1. Map of Upper Cook Inlet, Alaska, showing location of Offshore Test Fish transect.






Figure 2. Daily (upper) and cumulative (lower) CPUE for sockeye salmon recorded during the 1991 offshore test fish project, Upper Cook Inlet, Alaska.



Cumulative proportions estimated for the sockeye salmon return to Upper Cook Inlet, Alaska, in 1991. Figure 3.



Cumulative proportions for all stations combined, station 6a only, and all stations except 6a for the sockeye salmon return to Upper Cook Inlet, Alaska, in 1991. Figure 4.

Appendix A.1. Summary of pink salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fishing project, 1991.

	Number	Mean Fishing	Cat	ch	· CP	UE
Date	of Stations	Time (min)	Daily	Cum	Daily	Cum
01 July 02 July 03 July 04 July 05 July 06 July 07 July 08 July 10 July 11 July 12 July 13 July 14 July 15 July 16 July 17 July 18 July 20 July 21 July 22 July 23 July 24 July 25 July 27 July 28 July 29 July 30 July	545555555555555555555555555555555555555	195.5 131.5 186.0 188.5 186.0 190.5 188.5 186.5 186.5 204.5 188.5 217.5 180.0 230.0 195.0 197.5 196.0 192.5 180.0 200.0 194.5 178.0 191.5 191.5 191.5 191.5 191.5 191.5	0 0 0 0 0 0 0 0 0 0 1 1 0 3 1 3 1 0 6 1 1 1 1 0 0 2 1	0 0 0 0 0 0 0 0 0 0 0 1 2 2 5 6 9 10 16 17 21 32 32 32 32 33 33 34 35 36 36 37 37 37 37 37 37 37 37 37 37 37 37 37	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0

Appendix A.2. Estimated pink salmon catch by date and station, Upper Cook Inlet offshore test fishing project, 1991.

		Stat	Station Number					
Date	4	5	6	7	8	Total		
01 July	0	0	0	0	0	0		
02 July	0	0	0	0		0		
03 July	0	0	0	0	0	0		
04 July	0	0	0	0	0	0		
05 July	0	0	0	0	0	0		
06 July 07 July	0	0 0	0 0	0 0	0	0		
07 July 08 July	0 0	0	0	0	0	0		
09 July	0	0	0	0	0	0		
10 July	0	1	Ö	0	0	1		
11 July	Ŏ	Ō	ĭ	ŏ	ŏ	i		
12 July	Ö	Ö	Ō	Ō	Ö	Ō		
13 July	1	0	2	0	. 0	3		
14 July	0	1	0	0	0	1		
15 July	0	0	3	0	0	3		
16 July	0	0	0	0	1	1		
17 July	0	0	0	0	0	0		
18 July	1	3	0	1	1	6		
19 July	1	0	0	0	0	1		
20 July 21 July	1	2 0	0	1 1	0	4		
21 July 22 July	0 0	1	1 0	0	0 1	2 2		
23 July	0	0	1	0	3	4		
24 July	0	ő	i	Ő	0	1		
25 July	Ŏ	Ŏ	Ō	Ŏ	ĭ	i		
26 July	Ō	Ö	ĺ	Ō	Ō	ī		
27 July	0	0	0	0	0	Ō		
28 July			0	0	0	0		
29 July	0	1	1	0	0	2		
30 July	0	0	0	0	I	1		
Total	4	9	11	3	8	35		
%	11.4	25.7	31.4	8.6	22.9	100.0		

Appendix A.3. Estimated pink salmon CPUE by date and station, Upper Cook Inlet offshore test fishing project, 1991.

		Stat	ion Number			
Date	4	5	6	7	8	Total
01 July 02 July 03 July 04 July 05 July 06 July 07 July 08 July 10 July 11 July 12 July 13 July 14 July 15 July 16 July 17 July 18 July 20 July 21 July 21 July 22 July 23 July 24 July 25 July 26 July 27 July 28 July 29 July 30 July	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.6 0.0 2.1 1.5 0.7 0.8 2.8 1.3 1.6 2.8 0.7 0.0 0.0
Total %	4 12.4	9 27.3	11 28.0	3 9.1	8 23.2	35 100.0

Appendix B.1. Summary of chum salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fishing project, 1991.

	Number	Mean Fishing	Cat	cch	C	PUE
Date	of Stations	Time (min)	Daily	Cum	Daily	Cum
01 July 02 July 03 July 04 July 05 July 06 July 07 July 09 July 10 July 11 July 12 July 13 July 14 July 15 July 17 July 18 July 19 July 20 July 21 July 22 July 23 July 24 July 25 July 26 July 27 July 28 July 29 July 30 July	54555555555555555555555555555555555555	195.5 131.5 186.0 188.5 186.0 190.5 188.5 186.5 186.5 204.5 188.5 217.5 180.0 230.0 195.0 187.5 196.0 192.5 180.0 200.0 200.0 194.5 178.0 191.5 191.5 191.5 191.5 191.5	2 1 0 3 2 1 8 8 5 9 15 1 32 7 44 11 3 14 9 12 24 46 24 3 19 64 77 64 161 2	2 3 3 6 8 9 17 25 30 54 55 87 9 18 149 156 177 217 228 444 508 669 671	1.5 0.8 0.0 2.4 1.6 0.8 6.3 3.9 9.5 7 19.3 21.9 8.2 2.3 10.8 12.4 15.5 29.1 17.3 2.5 13.9 41.6 44.0 38.4 101.1 1.7	1.5 2.3 4.7 6.3 7.1 13.6 23.6 30.5 39.9 40.7 605.8 87.7 98.2 108.3 115.1 127.5 143.0 205.5 192.5 192.5 291.5 291.5 291.5 291.5 291.7 432.7

Appendix B.2. Estimated chum salmon catch by date and station, Upper Cook Inlet offshore test fishing project, 1991.

		St	ation Numb	per		
Date	4	, 5	6	7	8	Total
01 July	1	0	1	0	0	2
02 July	0	0	1	0	-	1
03 July	0	0	0	0	0	0
04 July	0 0	2 2	0	1	0	3 2
05 July 06 July	0	0	0 1	0 0	0 0	1
07 July	0	0	8	0	0	
08 July	Ŏ	ŏ	3	5	Ö	8
09 July	2	2	ĺ	Õ	ő	5
10 July	0	2 8	Ō	Ö	ĺ	8 8 5 9
11 July	0	9	6	0	0	15
12 July	0	1	0	0	. 0	1
13 July	0	17	4	11	0	32
14 July	0	0	2	4	1	7
15 July	0	1	41	2 6	0	44
16 July	0	0	4		1	11
17 July 18 July	1 2	0	2 6	0 1	0 0	3 14
19 July	1	5 3 6	5	0	0	9
20 July	3	6	2	Ö	1	12
21 July	Ō	3	8	12	î	24
22 July	0	0	8	35	3	46
23 July	1	0	6	5	12	24
24 July	2	0	0	1	0	3
25 July	0	0	0	10	9	19
26 July	0	1	62	1	0	64
27 July	0	0	60	15	2	77
28 July 29 July	1	63	64 35	0 56	0	64
30 July	0	0	35 2	0	6 0	161 2
Total	14	123	332	165	37	671
%	2.1	18.3	49.5	24.6	5.5	100.0

Appendix B.3. Estimated chum salmon CPUE by date and station, Upper Cook Inlet offshore test fishing project, 1991.

		Stat	ion Number			
Date	4	5	6	7	. 8	Total
01 July 02 July 03 July 04 July 05 July 06 July 07 July 08 July 10 July 11 July 12 July 13 July 14 July 15 July 16 July 17 July 18 July 19 July 20 July 21 July 21 July 22 July 23 July 24 July 25 July 26 July 27 July 28 July 29 July	0.7 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.0 1.5 1.6 0.0 0.0 0.0 1.5 6.1 5.6 0.7 10.7 0.0 0.8 0.0 0.0 3.3 2.2 3.5 2.4 0.0 0.0 0.0 0.0	0.8 0.8 0.0 0.0 0.0 0.8 6.2 2.3 0.0 3.9 0.7 1.4 20.0 3.2 1.5 4.3 3.8 5.7 4.5 6.7 0.0 4.7 0.0 4.7 0.0 4.7 0.0 4.7 1.7	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	1.5 0.8 0.0 2.4 1.6 0.8 6.2 6.3 3.9 6.9 9.5 0.7 19.9 5.3 21.9 8.2 2.3 10.1 6.8 12.4 15.5 29.1 17.3 2.5 13.9 41.6 44.0 38.4 101.1
Total	11.6	79.1	209.9	105.2	26.9	432.7
%	2.7	18.3	48.5	24.3	6.2	100.0

Appendix C.1. Summary of coho salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fishing project, 1991.

	Number	Mean Fishing	Cat	ch	CPU	JE
Date	of Stations	Time (min)	Daily	Cum	Daily	Cum
01 July 02 July 03 July 04 July 05 July 06 July 07 July 08 July 10 July 11 July 12 July 13 July 14 July 15 July 16 July 17 July 18 July 19 July 20 July 21 July 21 July 22 July 23 July 24 July 25 July 26 July 27 July 28 July 29 July 30 July	545555555555555555555555555555555555555	195.5 131.5 186.0 188.5 186.0 190.5 188.5 186.5 189.5 186.5 204.5 188.5 217.5 180.0 230.0 195.0 187.5 180.0 200.0 200.0 200.0 194.5 178.0 191.5 192.5 191.5 191.5 191.5	0 0 2 2 0 2 1 0 6 7 28 5 42 44 11 24 15 10 20 41 89 42 30 41 89 42 30 41 53 42 41 53 64 71 53 64 71 53 64 71 71 71 71 71 71 71 71 71 71 71 71 71	0 0 2 4 4 6 7 7 13 20 48 53 95 139 150 174 189 199 260 349 391 425 444 478 549 602 689 690	0.0 0.0 1.6 1.6 0.0 1.5 0.8 0.0 4.7 5.5 18.3 3.8 26.3 35.8 17.9 11.2 7.8 15.2 37.6 59.9 28.5 21.7 3.4 14.2 23.1 45.4 35.5 8 0.8	0.0 0.0 1.6 3.2 3.2 4.7 5.5 5.5 10.2 15.7 34.0 37.8 64.1 97.1 102.9 120.8 132.0 139.8 155.0 192.6 252.5 281.0 302.7 306.1 320.3 343.4 388.8 424.3 481.1 481.9

Appendix C.3. Estimated coho salmon CPUE by date and station, Upper Cook Inlet offshore test fishing project, 1991.

		S-	tation Num	ber		
Date	4	5	6	7	8	Total
01 July 02 July 03 July 04 July 05 July 06 July 07 July 08 July 10 July 11 July 12 July 13 July 14 July 15 July 16 July 17 July 18 July 19 July 20 July 21 July 22 July 23 July 24 July 25 July 26 July 27 July 28 July 29 July 30 July	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.8 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.0 1.6 0.0 1.5 0.0 0.0 2.4 12.9 0.8 8.9 11.4 11.4 16.0 2.3 0.8 3.3 17.4 13.9 25.2 11.0 0.0	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 13.4 18.2 2.1 7.1 0.0 4.1 0.0 1.7 39.8 13.7 7.0 0.0 3.0 2.4 26.6 3.7 19.4 0.0	0.0 0.8 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 1.6 0.0 1.5 0.0 4.7 5.3 8.3 33.8 26.3 5.8 17.9 11.2 8.5 17.2 8.5 17.2 8.5 17.2 8.5 17.2 8.5 17.2 8.5 14.2 23.1 45.4 14.2 24.1 45.5 16.0 16.0 16.0 16.0 16.0 16.0 16.0 16.0
Total	17.9	58.0	177.4	164.5	64.1	481.9
%	3.7	12.0	36.8	34.1	13.3	100.0

Appendix D.2. Estimated chinook salmon catch by date and station, Upper Cook Inlet offshore test fishing project, 1991.

		Station Number					
Date	4	5	6	7	8	Tota	
01 July	0	0	. 0	0	0	0	
02 July	0	0	0	0	•	0	
03 July	0	0	0	0 .	0	0	
04 July	0	0	0	0	0	0	
05 July	0	0	0	0	0	0	
o6 July	0	0	0	0	0	0	
o7 July	0	0	1	1	0	2	
08 July	0	0	0	0	0	0	
09 July	0	0	0	0	0	0	
10 July	0	0	0	0	0	0	
11 July	0	0	0	0	0	0	
12 July	0	0	0	0	: 0	0	
13 July	0	0	0	0	0	0	
14 July	0	0	0	0	0 0	0	
15 July	0	0 0	0	0	0	0 0	
16 July	0	0	0	0	0	0	
17 July	0	0	0 0	0 0	0	0	
18 July 19 July	0 0	0	0	0	0	0	
19 July 20 July	0	0	0	0	0	0	
20 July 21 July	0	0	0	0	0	0	
22 July	0	0	0	Ö	Ö	0	
23 July	0	Ö	0	Ö	Ö	0	
24. July	0	Ö	ő	ŏ	Ö	Ö	
25 July	Ö	Ö	ő	ŏ	Ŏ	Ō	
26 July	Ŏ	Ö	Ŏ	Ö	Ö	Ö	
27 July	Ö	Ö	Ŏ	Ö	Ö	Ö	
28 July	•	·	Ö	Ŏ	Ö	Ö	
29 July	0	0	Ö	Ŏ	Ö	Ö	
30 July	0	0	Ö	Ō	0	Ō	
Total	0	0	1	1	0	2	
% .	0.0	0.0	50.0	50.0	0.0	100.0	

Appendix D.3. Estimated chinook salmon CPUE by date and station, Upper Cook Inlet offshore test fishing project, 1991.

		St				
Date	4	5	6	7	. 8	Total
01 July 02 July 03 July 04 July 05 July 06 July 07 July 09 July 10 July 11 July 12 July 13 July 14 July 15 July 16 July 17 July 20 July 21 July 21 July 22 July 23 July 24 July 25 July 26 July 27 July 28 July 29 July 30 July			0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0		0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0
Total	0.0	0.0	0.8	0.8	0.0	1.6
%	0.0	0.0	50.0	50.0	0.0	100.0

Appendix E.1. Entry pattern of sockeye salmon into Upper Cook Inlet, Alaska, 1991, estimated from daily CPUE measured at the latitude of Anchor Point.

Appendix F.1. Environmental measurements and observations made in Upper Cook Inlet, 1991, Alaska during the offshore test fishing project.

Date	Station	Air Temp (c)	Water Temp (c)	Wind Vel (knots)	Wind Dir	Tide Stage	Water Depth (f)	Secchi (m)
Ol July	4 5 6 7	8 10 11 11	8.5 10.0 10.0 10.0	6 3 8 10	5 5 5 6	3 2 4 4	25.5 38.5 45.5 45.5	6.0 2.5 3.0 3.0
02 July	8 7 6 5 4	10 9 9 10	10.0 10.0 9.0 9.0	10 3 5 1 3	6 5 4 4	4 4 3 3	29.5 46.5 46.5 38.0 25.0	1.5 3.5 3.0 4.5 9.0
03 July	4 5 6 7	11 10 12 13 10	8.0 9.0 10.0 10.0	15 17 20 19	4 4 5 5 5 5 5	3 3 3 3 3 3 2 4	24.5 35.5 46.5 45.0	8.0 3.0 2.5 2.0
04 July	8 8 7 6 5	11 8 9 8 9	11.0 10.0 10.0 10.0 9.5	18 16 15 17 19	4 4 4	2 4 4 4 1	30.0 29.0 46.0 47.0 38.0	2.0 2.0 3.0 3.0 3.5
05 July	4 4 5 6 7	9 9 10 10 11	8.5 9.0 10.0 11.0 10.0	17 1 8 4	556666	1 1	26.0 26.5 37.0 48.0 44.0	4.0 7.0 3.0 3.0 3.0
06 July	, 8 8 7 6 5	12 10 11 10	10.0 11.0 10.0 10.0	8 2 0 1 2 3 2	6 0 1 1	3 3 3 3 2 4 4	29.5 29.5 45.0 47.0 39.0	3.0 2.0 3.0 4.0 6.0
07 July	4 4 5 6 7	10 10 11 13 14	10.0 8.0 8.5 9.0 9.5	2 0 0 0	2 0 0 0 6	4 4 4 4	26.0 27.0 37.5 50.0 46.0	11.5 11.0 8.0 3.5 3.0
08 July	8 8 7 6 5	12 9 9 9	9.0 9.0 9.5 10.0 8.0	3 10 12 10 10 7	5 4 4 4 4 5	3 3 3 3 2 4	30.0 29.0 45.0 44.0 37.0 27.0	3.0 3.0 3.5 2.5 5.0 6.0

Appendix F.1. (p 2 of 4)

Date	Station	Air Temp (c)	Water Temp (c)	Wind Vel (knots)	Wind Dir	Tide Stage o	Water Depth (f)	Secchi (m)
09 July	4 5 6 7	9 9 11 10	9.0 9.0 8.0 10.0	13 10 7 5	2 1 1 5	4 4 4 4	26.0 38.0 49.0 45.0	8.0 10.0 6.0 3.0
10 July	8 8 7 6	11 9 9 9	9.0 10.0 10.0 10.0 8.5	5 5 11 13 9 6	1 5 5 5 5 5	1 3 3 3 3 2 2	27.0 29.0 43.5 45.0 38.5	3.5 3.5 4.0 2.5 6.0
11 July	5 4 4 5 6 7	10 9 13 11	11.0 11.0 8.0 8.5	0 0 0 5	0 0 0 2 0	2 2 4 4 4	25.5 27.0 37.5 47.0 47.0	5.0 8.0 6.5 5.0 3.5
12 July	8 8 7	10 11 10 9 9	9.0 9.0 9.0 9.0	1 0 3 0	8 0 5 0	3 3 3 3 3 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3	29.0 31.5 45.0 46.5 40.0	2.0 3.5 3.0 3.0 4.0
13 July	6 5 4 4 5 6 7	10 10 10 10	8.5 9.0 9.0 10.0 9.0	0 0 5 12 7	0 0 5 6 5	3 3 4 4	25.0 25.0 40.0 48.0	5.0 4.0 4.0 2.5
14 July	8 8 7	10 11 9 10 10	10.0 9.0 10.0 9.0 10.0	9 8 9 8 11	056564545554	4 4	50.0 28.0 28.0 51.0 45.0	4.0 3.0 2.0 3.0 3.0
15 July	6 5 4 5 6	11 11 12 13 14	9.0 8.0 8.0 10.0	7 7 6 7 4	4	1 3 3 3 3 3 4	35.0 26.0 24.0 36.0 45.0	4.5 6.0 7.0 3.5 2.5
16 July	7 8 8 7 6 5	13 11 9 10 9	11.0 10.0 10.0 10.0 10.0	6 3 5 6 2 3 2	6 4 2 2 2 2	4 4 3 3 3 3	43.5 26.0 31.0 46.0 49.0	2.0 3.0 3.5 3.0
17 July	5 4 4	10 9 9	9.0 9.0 9.0	2 10	2 2 5	3 3 1	41.0 26.0 26.0	4.0 6.0 6.5

Appendix F.1. (p. 3 of 4)

Date	Station	Air Temp (c)	Water Temp (c)	Wind Vel (knots)	Wind Dir	Tide Stage⁵	Water Depth (f)	Secchi (m)
17 July	5 6 7	9 10 11	9.0 10.0 10.0	12 17 16	6 6 5	3 3 2	38.0 48.0 45.0	4.0 3.0 3.0
18 July	8 8 7 6 5	13 10 12 11 10	10.0 11.0 10.0 10.0 10.0	18 5 3 4 10	4 1 1 1	4 4 1	25.0 32.0 46.0 45.0 44.0	1.5 2.0 3.0 3.0 3.5
19 July	5 4 4 5 6 7	10 10 11 14 15	9.0 9.0 11.0 10.0	10 7 0 0	6 5 0 0	3 4 1 3 3	26.0 28.0 38.0 49.0 45.0	5.5 5.5 4.0 3.0 3.5
20 July	, 8 8 7 6 5	12 10 11 11	10.0 10.0 11.0 10.5 10.0	2 0 0 1 4	4 0 0 8 4	3 4 4 4 4 4	28.0 27.0 43.0 43.0 37.0	3.0 3.0 2.5 3.0 4.0
21 July	4 4 5 6 7	10 11 13 15	9.0 9.0 10.0 11.0 12.0	3 0 0 0	4 0 0 0 7	4 4 4 4	28.0 28.0 33.0 50.0 44.0	4.5 6.5 5.0 2.5 3.0
22 July	8 8 7 6 5	16 11 13 13	11.0 11.0 11.5 11.5	0 0 2 5	, 0 5 6 0	3 3 4 4 4	30.0 30.0 45.0 45.0 37.0	3.0 3.0 2.5 3.0 5.0
23-Jul	4 4 5 6 7	13 9 10 12 13	9.0 9.0 10.0 11.0	0 18 13 15	0 8 1 1 2	4 4 4 4 4	27.0 25.0 46.0 47.0 46.0	6.0 4.5 4.5 4.0 3.0
24 July	, 8 8 7 6 5	14 10 11 11	11.5 11.5 11.5 11.0 10.0	13 12 18 19 23	2 8 1 2 8	4 3 3 3 2	28.0 31.0 45.0 45.0 37.0	2.5 3.5 3.5 4.0 4.0
25 July	4 4 5	10 14 13	11.0 10.0 10.0	20 0 0	8 0 0	4 3 2	28.0 25.0 42.0	5.5 7.0 7.5

-Continued-

Appendix F.1. (p 4 of 4)

		Air	Water	Wind			Water	
Date	Station	Temp (c)	Temp (c)	Vel (knots)	Wind Dir	Tide Stage	Depth (f)	Secchi (m)
25 11.4		13	10.0	0	0	. 4	47.0	5.0
25 July	6 7	15	10.0	0 0	0	4	45.0	3.5
	8	13	11.0	ő	Ö	4	32.0	3.0
26 July	8 7	10	11.0	4	5	3	29.5	3.0
		12	11.0	7	4	3	42.5	3.0
	6	12	11.0	8	4	3	46.0	3.0
	6 5 4	14	11.0	7	4	3 2	40.0 28.5	4.0 10.0
27 July		14 13	10.0 9.0	0 7	0 4	3	24.5	6.5
Z/ July	4 5	13	10.0	8	4	3 3 3 2 3 3	37.0	5.0
	6	17	11.0	4	4	2	46.0	3.5
	7	14	10.0	3		4	36.0	2.5
	8 8 7	13	11.5	3 4 2	4 4 5 5	4	26.0	2.5
28 July	8	13	10.0	2	5	3	31.0 44.0	3.0 2.5
		11 13	11.0 11.0	12 15	5 5	3	44.0	3.5
29 July	6 4	13	9.0	15	6	4 3 3 3 3 3	25.0	6.5
25 outy	5	12	11.0	18	6	3	36.0	2.5
	4 5 6	13	11.5	18	6	4	46.0	2.5
	7	11	11.0	16	6	4	47.0	2.5
	8 8	10	11.0	11	6	4	31.0	2.5
	8	9	11.0	4	1	1	27.0	3.0
	7 6	10 9	11.0 10.0	0 0	0 0	- 3 3	51.0 47.0	3.5 3.5
	5 5	10	10.0	0	0	3	37.0	5.0
	6 5 4	10	10.0	Ö	0	3 3	26.0	8.0